NOAA Climate Science & Services Monthly Climate Update





Jake Crouch

Climate scientist, climate monitoring branch, NOAA
National Centers for Environmental Information

William (Billy) Sweet

Oceanographer, NOAA Center for Oceanographic Products and Services


Brad Pugh

Meteorologist & Seasonal Forecaster, NOAA Climate Prediction Center, Operations Prediction Branch

Contiguous U.S. Temperatures 2017


May was slightly warmer than average

Temperature Percentiles May 2017 Period: 1895-2017 (123 years)




Spring was the 8th warmest on record

Temperature Percentiles Spring 2017 Period: 1895-2017 (123 years)



Jan-May was 2nd warmest on record

Temperature Percentiles Jan-May 2017 Period: 1895-2017 (123 years)



- Warmth along the West Coast and in Florida
- Below-average temperatures in the south-central US
- Warm nighttime temperatures and cool daytime temperatures in the Midwest to Northeast
- Above-average temperatures in the Rockies, South, and Southeast
- Much of the spring warmth came early in the season – 11th warmest April and 9th warmest March
- Record warm in the Southern Plains and Southeast: FL, GA, NC, SC, and TX record warm
- Below-average temperatures in parts of the Northwest. WA only cooler than average state






Contiguous U.S. Precipitation 2017


May was wetter than average

Precipitation Percentiles May 2017 Period: 1895-2017 (123 years)




Spring was the 11th wettest on record

Precipitation Percentiles Spring 2017 Period: 1895-2017 (123 years)



Jan-May was 4th wettest on record

Precipitation Percentiles Jan-May 2017 Period: 1895-2017 (123 years)



- Wetter than average in the East

 record precipitation for parts of the central Appalachians
- Drier than average in the Northern and Southern Plains.
 Drought expanded in the Northern Plains
- Wetter than average in the Northwest
- Record precipitation in the Mid-Mississippi Valley and central Appalachians
- Drier than average in the Northern Plains, Southwest, and Florida

- Record and near-record precipitation across much of the West.
- Wetter than average across the Central Plains to Midwest and Northeast
- Drier than average in the Northern Plains and Florida




Current U.S. Drought


7.3% of Contiguous U.S. in Drought

2.6 percentage points since early Apr)

- Improvement: Central and Southern Plains, Southeast, and Northeast
- Degradation: Northern Plains and Southwest
- Outside CONUS: Drought improved in Hawaii with abnormally dry conditions in Alaska








High Tide Flooding in U.S.

- Due to sea level rise (SLR), flood frequency trends (linear, quadratic) are rapidly increasing
- Upwards of 300 to 1000% increase since 1960s
- Elevation thresholds when impacts occur vary by location
- Seasonal high tides and minor wind events now cause high tide flooding in many locations






(Sweet et al., 2017)

High Tide Flooding and El Nino in U.S.


- During El Nino, West and East Coast regions typically (statistically significant) experience more tidal flooding than (trend) normal
- Models suggest a mild El Nino might form this year -- more flooding (red star)





2016 High Tide Flooding in U.S.


- High tide flooding broke 2 records (Charleston: 50 days; Savannah: 38 days) and tied 1 record (Key West: 14 days)
- Flooding in 2016 was about or above trend-expected values at most locations (130% higher on average compared to 1995)
- Flooding most frequent where 2016 sea levels higher and flood elevations lower




June 2017


2017 High Tide Flood 'Outlook'

Based upon trend (and % increase from El Nino pattern)



Sea Surface Temperatures and ENSO





SON OND

Time Period

NDJ

ASO

MJJ

2017


Sea surface temperatures

- Above normal SSTs in the central and eastern Pacific although the upperocean heat anomaly decreased this past month
- Above normal SSTs in the tropical Atlantic
- ENSO neutral conditions are present, given the state of the ocean and atmosphere

ENSO forecast

- ENSO neutral is the most likely outcome through the fall and winter 2017-18
- The chances of El Niño development are similar to climatology by the fall
- A repeat of La Niña is unlikely in 2017


Monthly Forecast (July)






Seasonal Forecast (July-Aug-Sep)

July-Aug-Sep Average Temperature Probability July-Aug-Sep Total Precipitation Probability



U.S. Drought Outlook





12

For More Information



www.ncdc.noaa.gov

TODAY'S PRESENTATION:

http://www.ncdc.noaa.gov/sotc/briefings

NOAA's National Centers for Environmental Information:

- Monthly climate reports (U.S. & Global): www.ncdc.noaa.gov/sotc/
- Dates for upcoming reports: http://www.ncdc.noaa.gov/monitoring-references/dyk/monthly-releases

NOAA's Climate Prediction Center: www.cpc.ncep.noaa.gov

U.S. Drought Monitor: http://drought.gov

Climate Portal: www.climate.gov

2016 State of High Tide Flooding Report:

https://www.ncdc.noaa.gov/sotc/national/2017/05/supplemental/page-1

NOAA Media Contacts:

- John.Leslie@noaa.gov , 301-713-0214 (NOAA Communications/NESDIS)
- Brady.Phillips@noaa.gov, 202-482-2365 (NOAA Communications/HQ)

